

Few-Body Correlations in Fermi Systems

M. Beyer

FB Physik, University of Rostock, 18051 Rostock, Germany

Interacting quantum systems with strong or long-range interactions exhibit quite a rich phase structure. Cluster formation and superconductivity are examples. These phenomena are also expected in the astrophysical context, e.g., during the formation or in the structure of neutron stars. To describe these phenomena a proper treatment has to go beyond the simple picture of noninteracting quasiparticles. An appealing formalism for a systematic approach is provided by the framework of Dyson equations. Within an equal (imaginary) time formalism Dyson equations can be derived for an arbitrary large cluster embedded in a medium [1]. For practical use and the sake of simplicity the medium is treated as uncorrelated to derive the respective n -body cluster Green functions. Further, we neglect “backward” propagating particles, so the Fock spaces for different number of particles n are disconnected. This way it is possible to derive effective in-medium n -body equations that can be solved rigorously with few-body techniques [2–13].

These resulting two-, three-, and four-body equations elaborated here include the dominant medium effects in a systematic way. These are the self energy corrections for masses and the Pauli blocking that in turn leads to a change of binding energies, viz. change of the masses of clusters, and change of reaction rates. Further, within this approach the critical temperatures for condensation (of bosons containing two or four particles) are calculated.

Defining $H_0 = \sum_{i=1}^n \varepsilon_i$ with the quasi-particle self energy

$$\varepsilon_1 = k_1^2/2m_1 + \sum_2 V_2(12, \widetilde{12})f_2 \quad (1)$$

and the Fermi function $f_1 \equiv f(\varepsilon_1) = 1/(e^{\beta(\varepsilon_1 - \mu)} + 1)$, the n -particle cluster resolvent G_0 is

$$G_0(z) = (z - H_0)^{-1} N \equiv R_0(z) N. \quad (2)$$

Here G_0 , H_0 , and N are matrices in n particle space and z denotes the Matsubara frequency [14]. The Pauli-blocking factors for n -particles are

$$N = \bar{f}_1 \bar{f}_2 \dots \bar{f}_n \pm f_1 f_2 \dots f_n, \quad \bar{f} = 1 - f \quad (3)$$

Note: $NR_0 = R_0N$. Defining the effective potential $V \equiv \sum_{\text{pairs } \alpha} N_2^\alpha V_2^\alpha$ the full and the channel resolvents are

$$G(z) = (z - H_0 - V)^{-1} N \equiv R(z)N, \quad (4)$$

$$G_\alpha(z) = (z - H_0 - N_2^\alpha V_2^\alpha)^{-1} N \equiv R_\alpha(z)N, \quad (5)$$

Note that $V^\dagger \neq V$ and $R(z)N \neq NR(z)$. For the scattering problem it is convenient to define the in-medium AGS operator $U_{\beta\alpha}(z)$

$$R(z) = \delta_{\alpha\beta} R_\beta(z) + R_\beta(z) U_{\beta\alpha}(z) R_\alpha(z) \quad (6)$$

that after some algebra leads to the in-medium AGS equation

$$U_{\beta\alpha}(z) = \bar{\delta}_{\beta\alpha} R_0(z)^{-1} + \sum_{\gamma} \bar{\delta}_{\beta\gamma} N_2^\gamma T_2^\gamma(z) R_0(z) U_{\gamma\alpha}(z), \quad (7)$$

where $\bar{\delta}_{\beta\alpha} = 1 - \delta_{\beta\alpha}$. The square of this t -operator is directly linked to the differential cross section for the scattering process $\alpha \rightarrow \beta$. The driving kernel consists of the two-body t -matrix derived in the same formalism, however given earlier and known as Feynman-Galitskii equation [14]

$$T_2^\gamma(z) = V_2^\gamma + V_2^\gamma N_2^\gamma R_0(z) T_2^\gamma(z).$$

A numerical solution using a coupled Yamaguchi potential has been given in Ref. [2]. For a temperature $T = 10$ MeV and the three-body system at rest in the medium results a given in Fig.1 For the bound state problem it is convenient to introduce form factors

$$|F_\beta\rangle = \sum_{\gamma} \bar{\delta}_{\beta\gamma} N_2^\gamma V_2^\gamma |\psi_{B_3}\rangle. \quad (8)$$

Since the potential is nonsymmetric right and left eigenvectors are different, although the bound state energies are the same,

$$|F_\alpha\rangle = \sum_{\beta} \bar{\delta}_{\alpha\beta} N_2^\beta T_2^\beta(B_3) R_0(B_3) |F_\beta\rangle, \quad (9)$$

$$|\tilde{F}_\alpha\rangle = \sum_{\beta} \bar{\delta}_{\alpha\beta} T_2^\beta(B_3) N_2^\beta R_0(B_3) |\tilde{F}_\beta\rangle. \quad (10)$$




FIG. 1. Break-up cross section for different densities of nuclear matter for temperature $T = 10$ MeV

The binding energy depends on μ , T , $P_{c.m.}$. Results for $P_{c.m.} = 0$ are given in Fig.2 for different potentials and temperatures. Note that the dependence on density is rather similar for two different potentials studied, although the binding energies for the isolated triton differs by 10% the Mott density is practically at the same place once the binding energies are renormalized to each other. For helium the Mott density is smaller due to the Coulomb force, however for asymmetric nuclear matter, e.g. $N_p/N_n \simeq 0.72$ (for the $^{129}\text{Xe} + ^{119}\text{Sn}$ reaction) this effect is compensated [15]. The dependence of the Mott effect on the momentum is given in Ref. [5].

We now turn to the four-body problem in matter. In addition to having different channels as for the three body system now the channels appear in different partitions that makes the four-body problem even more involved. The partitions of the four-body clusters are denoted by $\rho, \tau, \sigma, \dots$, e.g., $\rho = (123)(4), (234)(1), \dots$ for $3 + 1$ -type partitions, or $\rho = (12)(43), (23)(41), \dots$ for $2 + 2$ -type partitions. The two-body sub-channels are denoted by pair indices $\alpha, \beta, \gamma, \dots$, e.g. pairs (12), (24), $\dots$. The two- and three-body t -matrices have to be defined with respect to the partitions that leads to additional indices. The four-body in-medium homogeneous AGS equation are defined for the form factors




FIG. 2. Triton binding energy as a function of nuclear matter density. Dashed-dot-dot $T = 10$ MeV, other $T = 20$ MeV.

$$|\mathcal{F}_\beta^\sigma\rangle = \sum_\tau \bar{\delta}_{\sigma\tau} \sum_\alpha \bar{\delta}_{\beta\alpha}^\tau R_0^{-1}(B_4) |\psi_\alpha\rangle, \quad (11)$$

where $\bar{\delta}_{\beta\alpha}^\tau = \bar{\delta}_{\beta\alpha}$, if $\beta, \alpha \subset \tau$ and $\bar{\delta}_{\beta\alpha}^\rho = 0$ otherwise and $|\psi_\alpha\rangle$ is the α -particle wave function. They read [8]

$$|\mathcal{F}_\beta^\sigma\rangle = \sum_{\tau\gamma} \bar{\delta}_{\sigma\tau} U_{\beta\gamma}^\tau(B_4) R_0(B_4) N_2^\gamma T_2^\gamma(B_4) R_0(B_4) |\mathcal{F}_\gamma^\tau\rangle, \quad (12)$$

where $\alpha \subset \sigma, \gamma \subset \tau$. A numerical solution of this equation is rather complex. In order to reduce computational time we introduce a energy dependent pole expansion (EDPE) that has been proven useful in many application involving the α -particle and is accurate enough for the present purpose. However, we have to generalize the original version of the EDPE because of different right and left eigenvectors. Details will be omitted here, see [8]

In the *two-body* sub-system the EDPE reads

$$\begin{aligned} T_\gamma(z) &\simeq \sum_n |\tilde{\Gamma}_{\gamma n}(z)\rangle t_{\gamma n}(z) \langle \Gamma_{\gamma n}(z)| \simeq \sum_n |\tilde{g}_{\gamma n}\rangle t_{\gamma n}(z) \langle g_{\gamma n}| \\ &= \sum_n N_2^\gamma |g_{\gamma n}\rangle t_{\gamma n}(z) \langle g_{\gamma n}|. \end{aligned} \quad (13)$$

where we have chosen a Yamaguchi ansatz for the form factors for simplicity. Inserting this ansatz into the Feynman-Galitskii equation determines the propagator $t_{\gamma n}(z)$. In the *three-body* sub-system the EDPE expansion reads

$$\langle g_{\beta m}(z) | R_0(z) U_{\beta\gamma}^\tau(z) R_0(z) | \tilde{g}_{\gamma n}(z) \rangle \simeq \sum_{t,\mu\nu} |\tilde{\Gamma}_{\beta m}^{\tau t,\mu}(z)\rangle t_{\mu\nu}^{\tau t}(z) \langle \Gamma_{\gamma n}^{\tau t,\nu}(z) |. \quad (14)$$

with the three-body EDPE functions

$$|\tilde{\Gamma}_{\beta m}^{\tau t,\mu}(z)\rangle = \langle g_{\alpha n} | R_0(z) | \tilde{g}_{\beta m} \rangle t_{\beta m}(B_3) |\tilde{\Gamma}_{\beta m}^{\tau t,\mu}\rangle, \quad (15)$$

that we get from solving the proper Sturmian equations

$$\eta_{t,\mu} |\tilde{\Gamma}_{\alpha n}^{\tau t,\mu}\rangle = \sum_{\beta m} \langle g_{\alpha n} | R_0(B_3) | \tilde{g}_{\beta m} \rangle t_{\beta m}(B_3) |\tilde{\Gamma}_{\beta m}^{\tau t,\mu}\rangle \quad (16)$$

$$\eta_{t,\mu} |\Gamma_{\alpha n}^{\tau t,\mu}\rangle = \sum_{\beta m} \langle \tilde{g}_{\alpha n} | R_0(B_3) | g_{\beta m} \rangle t_{\beta m}(B_3) |\Gamma_{\beta m}^{\tau t,\mu}\rangle \quad (17)$$

Inserting everything into the homogeneous AGS equations allows us to redefine the form factors that are now operators in the coordinates of the $2+2$ or $3+1$ system, respectively

$$|\Gamma_\mu^{\sigma s}\rangle = \sum_{\beta m} \langle \Gamma_{\beta m,\nu}^{\sigma s}(B_4) | t_{\beta m}(B_4) \langle g_{\beta m}(B_4) | R_0(B_4) | \mathcal{F}_\beta^\sigma \rangle \quad (18)$$

and therefore the final homogeneous equation

$$|\Gamma_\mu^{\sigma s}\rangle = \sum_{\tau t} \sum_{\nu\kappa} \sum_{\gamma n} \bar{\delta}_{\sigma\tau} \langle \Gamma_{\gamma n}^{\sigma s,\nu}(B_4) | t_{\gamma n}(B_4) | \tilde{\Gamma}_{\gamma n}^{\sigma s,\mu}(B_4) \rangle t_{\mu\kappa}^{\tau t}(B_4) |\Gamma_\kappa^{\tau t}\rangle, \quad (19)$$

is an effective one-body equation with an effective potential $\mathcal{V}$ and an effective resolvent $\mathcal{G}_0$:

$$\mathcal{V}_{\mu\nu}^{\sigma s,\tau t}(z) = \sum_{\gamma n} \bar{\delta}_{\sigma\tau} \langle \Gamma_{\gamma n}^{\sigma s,\mu}(z) | t_{\gamma n}(z) | \tilde{\Gamma}_{\gamma n}^{\sigma s,\nu}(z) \rangle, \quad (20)$$

$$\mathcal{G}_{\mu\nu,0}^{\sigma s,\tau t}(z) = t_{\mu\nu}^{\tau t}(z). \quad (21)$$




FIG. 3. Energy dependence of the binding energy of the α -particle

The binding energies of the two-, three-, and four-body systems are shown in Fig.3 for a temperature of $T = 10$ MeV and a c.m. momentum of the respective cluster of $P_{\text{c.m.}} = 0$. The $B = 0$ line reflects the respective continuum. Investigating the zeros of the two-body Joost function the quasi deuteron survives as an anti-bound state (not resonance) with increasing densities, viz. for energies above the continuum [16]. The fate of the triton and of the α particle for $B > 0$ still needs to be investigated as well as a possible appearance of Efimov states related to $B \rightarrow 0$ of the sub-system. Since the Efimov states are 'excited' states, e.g. for the three-body system close to the $2 + 1$ threshold, their blocking may be smaller since the wave functions contain higher momentum components. Note that the slope of the binding energies as a function of densities for the larger clusters is also larger. This is a clear indication that the masses of the clusters change with increasing density. The Mott density of the α particle appears at

$$B_4(n_{\text{Mott}} = 0.19 \text{ fm}^{-3}, T = 10 \text{ MeV}, P_{\text{c.m.}} = 0) = 0$$

For comparison a perturbative result using Gaussian functions fitted to the charge radius of the α -particle is also given.




FIG. 4. Temperature vs. effective chemical potential ($\mu^* = \mu - \Sigma(0)$), $\Sigma(0)$ Hartree-Fock shift of the single particle energy at zero momentum. Lines show the critical temperature for pairing (solid) and quartetting. The dashed line shows a result given in [17], the diamonds show the solution of the AGS equation given in (19)

Finally, I address the question of a possible four-particle condensate or quartetting [17]. The condition is $B_4(n, T_c, P_{c.m.} = 0) = 4\mu$. From Fig.3 we argue that α condensation is likely, i.e.

$$T_c^\alpha > T_c^{NN},$$

where the critical temperature for α condensation turns out to be higher than for the pairing. However, for $\mu > 0$ the situation seems not so clear, since the four-body AGS equation (19) develops poles related to zeros in $4\mu - H_0$. Unlike the two-body case where these poles disappear because the numerator becomes as well zero at 2μ and T_c (viz. $1 - f_1 - f_2 \rightarrow 0$). A vanishing numerator is not obvious for the four-body case because there are more channels involved. It remains to be clarified, if the rapid fall of the critical temperature found in Ref. [17] using variational treatment with square-integrable functions remains, if one uses an exact treatment of the four-body problem. This is currently investigated.

Acknowledgment I am very grateful to S. Mattiello, C. Kuhrts, G. Röpke, P. Schuck, S.A. Sofianos, and W. Schadow, who have contributed in certain stages to some results presented here, and to T. Frederico for lively discussions. I gratefully acknowledge the warm and pleasant atmosphere at the Department of Physics during stays at UNISA, Pretoria. Work supported by Deutsche Forschungsgemeinschaft and University of South Africa.

-
- [1] J. Dukelsky, G. Röpke, P. Schuck, Nucl. Phys. A **628**, 17 (1998).
 - [2] M. Beyer, G. Ropke and A. Sedrakian, Phys. Lett. **B376** (1996) 7 [nucl-th/9601038].
 - [3] M. Beyer and G. Ropke, nucl-th/9706021.
 - [4] M. Beyer, Few Body Syst. Suppl. **10** (1999) 179 [nucl-th/9809002].
 - [5] M. Beyer, W. Schadow, C. Kuhrts and G. Ropke, Phys. Rev. C **60** (1999) 034004 [nucl-th/9902074].
 - [6] C. Kuhrts, M. Beyer and G. Ropke, Nucl. Phys. **A668** (2000) 137 [nucl-th/9908032].
 - [7] M. Beyer, C. Kuhrts, G. Ropke and P. D. Danielewicz, nucl-th/9910058.
 - [8] M. Beyer, S. A. Sofianos, C. Kuhrts, G. Ropke and P. Schuck, Phys. Lett. **B488** (2000) 247 [nucl-th/0003071].
 - [9] C. Kuhrts, M. Beyer, P. Danielewicz and G. Ropke, nucl-th/0009037.
 - [10] E.O. Alt, P. Grassberger, W. Sandhas, Nucl. Phys. **B 2** (1967) 167.
 - [11] W. Sandhas, Acta Physica Austriaca, Suppl. **XIII**, 679 (1974).
 - [12] E.O. Alt, P. Grassberger and W. Sandhas, Report E4-6688, JINR, Dunba 1972 and in *Few particle problems in the nuclear interaction* eds. I. Slaus et al. (North Holland, Amsterdam 1972) p. 299.
 - [13] W. Sandhas, Czech. J. Phys. **B 25**, 251 (1975).
 - [14] L.P. Kadanoff, G. Baym, *Quantum Theory of Many-Particle Systems* (Mc Graw-Hill, New York, 1962); A.L. Fetter, J.D. Walecka, *Quantum Theory of Many-Particle Systems*, (McGraw-Hill, New York, 1971).
 - [15] S. Mattiello, diploma thesis, 2000 (unpublished)
 - [16] M. Beyer and S.A. Sofianos, in preparation.
 - [17] G. Röpke, A. Schnell, P. Schuck, P. Nozière, Phys. Rev. Lett. **80** (1998) 3177